

Protoneutron star evolution and the neutrino driven wind in general relativistic neutrino radiation hydrodynamics simulations

T. Fischer^{*,†}, S. C. Whitehouse^{*}, A. Mezzacappa[‡], F.-K. Thielemann^{*,§} and M. Liebendörfer^{*}

March 2, 2010

Abstract

Massive stars end their life in an explosion event with kinetic energies of the order 10^{51} erg. Immediately after the explosion has been launched, a region of low density and high entropy forms behind the ejecta which is continuously subject to neutrino heating. The neutrinos emitted from the remnant at the center, the protoneutron star (PNS), heat the material above the PNS surface. This heat is partly converted into kinetic energy and the material accelerates to an outflow that is known as the neutrino driven wind. For the first time, we simulate the collapse, bounce, explosion and the neutrino driven wind phases consistently over more than 20 seconds. Our numerical model is based on spherically symmetric general relativistic radiation hydrodynamics using spectral three flavor Boltzmann neutrino transport. In simulations where no explosions are obtained naturally, we model neutrino driven explosions for low and intermediate mass Fe-core progenitor stars by enhancing the charged current reaction rates. In the case of a special progenitor star, the $8.8 M_{\odot}$ O-Ne-Mg-core, the explosion in spherical symmetry was obtained without enhanced opacities. The post explosion evolution is in qualitative agreement with static steady-state and parametrized dynamic models of the neutrino driven wind. On the other hand, we find generally smaller neutrino luminosities and mean neutrino energies as well as a different evolutionary behavior of the neutrino luminosities and mean neutrino energies. The neutrino driven wind is proton-rich for more than 10 seconds and the contraction of the PNS differs from the assumptions made for the conditions at the inner boundary in previous neutrino driven wind studies. Despite the moderately large entropies of about $100 k_B/\text{baryon}$ and the fast expansion timescale, the conditions found in our model are unlikely to favor r -process nucleosynthesis. The simulations are carried out until the neutrino driven wind settles down to a quasi-stationary state. After about 5 seconds post bounce, the peak temperature inside the PNS already starts to decrease due to the continued deleptonization. This moment determines the beginning of a cooling phase dominated by the emission of neutrinos. We discuss the physical conditions of the quasi-static PNS evolution and take the effects of deleptonization and mass accretion from early fallback into account.

Keywords: radiation hydrodynamics – Boltzmann neutrino transport – core collapse supernovae – neutrino driven explosions – neutrino driven wind – protoneutron stars

1 Introduction

Stars more massive than $8 M_{\odot}$ produce extended Fe-cores at the end of stellar evolution, which run into gravitational collapse due to pressure loss via photodisintegration and electron captures. The collapse halts at nuclear density, typically $2 - 4 \times 10^{14} \text{ g/cm}^3$ depending on the equation of state (EoS). However, the supersonically infalling material from the outer core continues to fall into the center. The core overshoots its equilibrium configuration and bounces back. A dynamic shock wave forms which propagates outwards and continuously loses energy due to the dissociation of heavy nuclei. As soon as the shock reaches the neutrinospheres, i.e. the neutrino energy and flavor dependent spheres of last scattering, additional electron captures emit a large amount of electron neutrinos. This burst of electron neutrinos, known as the deleptonization burst, carries away energy of several 10^{53} erg/s on a timescale of $10 - 20 \text{ ms}$.

^{*}Department of Physics, University of Basel, Klingelbergstrasse 82, 4056 Basel, Switzerland

[†]tobias.fischer@unibas.ch

[‡]Physics Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831-1200

[§]GSI Helmholtzzentrum für Schwerionenforschung GmbH, Planckstrasse 1, 64291 Darmstadt, Germany

This energy loss turns the expanding shock into a standing accretion shock (SAS) already about 5 ms after bounce. Due to the large energy in the neutrino radiation field, neutrino heating between the neutrinospheres and the SAS has long been investigated as a possible source to revive the SAS and trigger neutrino driven explosions (Bethe and Wilson (1985), Janka (2001), Janka et al. (2005), Mezzacappa et al. (2006)).

Up to now, neutrino driven explosions in spherical symmetry have only been obtained for the low mass $8.8 M_{\odot}$ O-Ne-Mg-core by Kitaura et al. (2006) and for low and intermediate mass progenitor stars by Sagert et al. (2009) assuming a hadron-quark phase transition during the early post bounce phase. On the other hand, multi-dimensional core collapse models with spectral neutrino transport have only recently become available and demonstrate the complexity of the underlying physical phenomena such as rotation and the development of fluid instabilities. These have been shown (see for example Miller et al. (1993), Herant et al. (1994), Burrows et al. (1995) and Janka and Müller (1996)) to increase the neutrino heating efficiency and help to understand aspherical explosions, see for example Bruenn et al. (2006) and Marek and Janka (2009). For a review on axially-symmetric neutrino driven explosions, see also Janka et al. (2008).

The following dynamical evolution of the PNS and hence the properties of the neutrino spectra emitted is determined by the mass accretion and the EoS. On a timescale of several seconds after the explosion has been launched, the region between the expanding explosion shock and the PNS at the center is subject to the formation of the neutrino driven wind as follows. Neutrinos continuously diffuse out of the hot PNS and heat the material on top of the PNS surface before they reach the neutrinospheres. We define the PNS surface to be the radius of the energy-integrated electron-neutrinosphere. The dominant neutrino heating contributions are given by the captures of electron-neutrinos and electron-antineutrinos on free nucleons. Matter is heated by neutrinos and the thermal energy is converted into kinetic energy, which accelerates material on top of the PNS surface to positive velocities. This matter outflow is known as the neutrino driven wind.

In this context, two particular studies are of special importance. The properties of the neutrino driven wind as described in Woosley et al. (1994) are based on the detailed radiation hydrodynamics simulation of a $20 M_{\odot}$ Fe-core progenitor applying the numerical model from Wilson and Mayle (1993). The simulation was carried until about 18 seconds after bounce. Another state-of-the-art model of that time was the explosion of the O-Ne-Mg-core by Mayle and Wilson (1988). Both investigations were milestones in the research of core collapse supernovae and are based on detailed neutrino input physics including neutrino transport, developed by J. R. Wilson. The results obtained, in particular the properties of the ejecta and the neutrino observables such as luminosities and energies, were considered the standard reference for more than 10 years. Neutrino driven wind studies used the results as parameters, where the conditions found indicated the possible site for the production of heavy elements via the r -process. In the simulations discussed in the present paper, we follow a similar approach as Wilson and Mayle (1993) and Mayle and Wilson (1988), where we apply the neutrino input physics based on Bruenn (1985). Although we find general agreement with the previous work, in particular the explosion phase, we find significant differences in the properties of the neutrino driven wind. The entropies per baryon are smaller by a factor of 2-3 and the wind stays proton-rich for more than 10 seconds for all our models. In addition, the neutrino luminosities and mean energies are generally smaller. The mean neutrino energies decrease with time, where they remain almost constant in the simulation of Woosley et al. (1994). The largest difference arises in the decreasing difference between the mean electron neutrino and antineutrino energies found in our simulations, i.e. the neutrino spectra become more similar with respect to time. The difference in the neutrino spectra in Woosley et al. (1994) remains large and even increases with time. Using the results from Woosley et al. (1994) as reference, Qian and Woosley (1996) analyzed the neutrino driven wind and formulated approximate analytical expressions for various properties of the neutrino driven wind, e.g. the neutrino heating rate, the electron fraction, the entropy per baryon and the mass outflow rate.

Based on the static wind equations, the results obtained in parameter studies (see e.g. Duncan et al. (1986), Hoffman et al. (1997), Thompson et al. (2001) and Thompson and Burrows (2001)) became known as static steady-state wind models, where Woosley and Baron (1992), Woosley et al. (1994), Takahashi et al. (1994) and Wittl et al. (1994) described the neutrino driven wind in a radiation-hydrodynamics context. Of special importance for the neutrino driven wind investigations is the impact on the nucleosynthesis. Most interesting is the possibility to explain the production of heavy elements via the r -process due to the large entropies, the fast expansion timescales and the low electron fraction of $Y_e < 0.5$ in the wind. Otsuki et al. (2000) explored general relativistic effects on the neutrino driven wind and investigated the possible impact on the nucleosynthesis. Recently, Wanajo (2006a) and Wanajo (2006b) investigated the neutrino driven wind with respect to the r - and rp -processes.

The possibility of supersonic wind velocities has been explored in most of the references. The supersonically expanding material in the wind collides with the much slower expanding and denser explosion ejecta. The material decelerates and a reverse shock forms which is known as the neutrino driven wind termination shock (first observed by Janka and Müller (1995) and Burrows et al. (1995)). Recently, Arcones et al. (2007) examined the post bounce phase of core collapse supernovae of several massive progenitor stars. Their models were launched from massive progenitor stars that were previously evolved through the core collapse, bounce and early post bounce phases using sophisticated radiation hydrodynamics based on spectral neutrino transport in spherical symmetry. The simulations are then continued applying a simplified radiation hydrodynamics description (see Scheck et al. (2006)), assuming large luminosities to trigger neutrino driven explosions in spherical symmetry. The neutrino driven wind develops supersonic outflow and the wind termination shock appears in all their models. Like most of the present neutrino driven wind studies, an interior boundary was assumed instead of simulating the PNS interior for the PNS contraction and the diffusion of neutrinos out of the PNS. However, steady-state wind studies could not predict the important dynamical features from the presence of the neutrino driven wind termination shock, especially the deceleration of the wind material and the consequent entropy as well as density and temperature increase during the deceleration. In this respect, the investigation from Arcones et al. (2007) was a milestone in modeling the neutrino driven wind consistently. On the other hand, they were focusing on parameters (luminosities and mean neutrino energies) in agreement with the simulations of Bethe and Wilson (1985) and Woosley et al. (1994). Although the dynamics is in general agreement, several properties of the neutrino driven wind as well as the neutrino spectra emitted differ significantly.

The present paper follows a different approach. We simulate consistently the dynamical evolution of the collapse, bounce and post bounce phases until the neutrino driven wind phase for more than 20 seconds. The simulations are launched from the $8.8 M_{\odot}$ O-Ne-Mg-core from Nomoto (1983,1984,1987) and the 10.8 and $18 M_{\odot}$ Fe-core progenitors from Woosley et al. (2002). Our numerical model is based on general relativistic radiation hydrodynamics with spectral three-flavor Boltzmann neutrino transport in spherical symmetry. The explosion mechanism of massive Fe-core progenitors is an active subject of research. To model neutrino driven explosions for such progenitors in spherical symmetry, we enhance the electronic charged current reaction rates artificially which increases the neutrino energy deposition and revives the SAS. The mechanism including the tuned neutrino reaction rates will be further discussed in §2 and §3. Such explosion models were investigated with respect to the nucleosynthesis by Fröhlich et al. (2006 a-c). Here, we report on the formation of the neutrino driven wind and the possibility of the wind developing supersonic velocities and hence the formation of the wind termination shock. In addition, we will also illustrate the explosion and the neutrino driven wind phases of the O-Ne-Mg core, where an explosion is found in spherical symmetry even without artificially enhanced neutrino opacities. The results are in qualitative agreement with those of Mayle and Wilson (1988) and Kitauro et al. (2006), who used a different EoS.

We find that for low progenitor masses, the neutrino driven wind termination shock will develop, using the tuned neutrino reaction rates. When the neutrino reaction rates are switched back to the standard opacities given in Bruenn (1985), the neutrino driven wind develops only a subsonic matter outflow. For intermediate progenitor masses, the neutrino driven wind remains subsonic even with the artificially enhanced neutrino emission and absorption rates. Since the neutrino driven wind depends sensitively on the emitted neutrino spectra at the neutrospheres, we believe accurate neutrino transport and general relativity in the presence of strong gravitational fields are essential in order to describe the dynamical evolution. Furthermore, the accurate modeling of the electron fraction in the wind is essential for nucleosynthesis calculations, which can only be obtained solving the neutrino transport equation. In addition, it is beyond the present computational capabilities to carry multi-dimensional simulations with neutrino transport to several seconds after bounce. Hence, our investigations are performed in spherical symmetry where we simulate the entire PNS interior rather than approximating an interior boundary. We find significant discrepancies in comparison with the assumptions made in previous wind studies. Material is found to be proton rich for more than 10 seconds, where most wind models assume luminosities and mean neutrino energies such that the neutrino driven wind becomes neutron-rich. We question the validity of the approximations made in such wind studies. We believe that an accurate and consistent modeling of the physical conditions in the wind is essential, especially in order to be able to draw conclusions with respect to nucleosynthesis.

The paper is organized as follows. In §2, we will present our spherically symmetric core collapse model. §3 is devoted to the explosion phase of neutrino driven explosions in spherical symmetry. We examine the $8.8 M_{\odot}$ progenitor model from Nomoto (1983,1984,1987) using the standard neutrino opacities and the 10.8 and $18 M_{\odot}$ progenitor models from Woosley et al. (2002) using artificially enhanced neutrino reaction rates. In §4 we discuss the conditions for the

formation of the neutrino driven wind and the possibility for the wind to develop supersonic velocities. We discuss in §5 the electron fraction approximation used in the literature. Since a generally neutron-rich neutrino driven wind is found in many previous and present wind studies, we illustrate the differences and investigate why we find a generally proton-rich wind. §6 is dedicated to the long term post bounce evolution for more than 20 seconds. In §7 we discuss the results and emphasize the main differences of the present investigation to previous wind studies. Finally we close with a summary in §8.

2 The model

Our core collapse model Agile-Boltztran is based on general relativistic radiation hydrodynamics in spherical symmetry, using three-flavor (anti)neutrino Boltzmann transport. For details see Mezzacappa and Bruenn (1993 a-c), Mezzacappa and Messer (1999), Liebendörfer et al. (2001a,b) and Liebendörfer et al. (2004). For this study we include the neutrino input physics based on Bruenn (1985). The charged current reactions considered

$$e^- + p \rightleftharpoons n + \nu_e, \quad (1)$$

$$e^+ + n \rightleftharpoons p + \bar{\nu}_e, \quad (2)$$

$$e^- + \langle A, Z \rangle \rightleftharpoons \langle A, Z-1 \rangle + \nu_e, \quad (3)$$

are electron and positron captures at free nucleons as well as electron captures at nuclei. The nuclei are characterized by an average atomic mass and charge $\langle A, Z \rangle$. In addition, the standard scattering reactions considered are iso-energetic neutrino nucleon ($N \in \{n, p\}$) and nuclei ($N = \langle A, Z \rangle$) scattering,

$$\nu + N \rightleftharpoons \nu + N,$$

where $\nu \in \{\nu_e, \nu_{\mu/\tau}\}$ (equivalent for antineutrinos $\bar{\nu}$), and neutrino electron/positron scattering

$$\nu + e^\pm \rightleftharpoons \nu + e^\pm.$$

The classical neutrino pair process is electron-positron annihilation,

$$e^- + e^+ \rightleftharpoons \nu + \bar{\nu}.$$

The standard neutrino energy E dependent emissivity $j(E)$ and absorptivity $\chi(E)$ for the charged current reactions as well as the scattering and pair-reaction kernels are given in Bruenn (1985) based on Yueh and Buchler (1976) and Schinder and Shapiro (1982). The additional pair-process nucleon-nucleon-Bremsstrahlung,

$$N + N \rightleftharpoons N + N + \nu + \bar{\nu},$$

has been implemented into our model according to Thompson and Burrows (2001) and is also taken into account. The emission of μ/τ -neutrino pairs via the annihilation of trapped electron-neutrino pairs,

$$\nu_e + \bar{\nu}_e \rightleftharpoons \nu_{\mu/\tau} + \bar{\nu}_{\mu/\tau},$$

as well as contributions from nucleon-recoil and weak magnetism as studied in Horowitz (2002) are investigated in Fischer et al. (2009) and are not taken into account in the present study of the neutrino driven wind.

2.1 Recent improvements of the adaptive grid

Long-term simulations of the supernova post bounce phase with Agile-Boltztran lead to a very large contrast of densities, reaching from $\sim 10^{15} \text{ g/cm}^3$ at the center of the protoneutron star (PNS) to densities of order g/cm^3 and smaller in the outer layers. The version of Agile described in Liebendörfer et al. (2002) is not able to resolve such large density contrasts. If the enclosed mass a is large and the density in one zone very small, then the evaluation of the mass contained in the zone according to Eq. (39) in Liebendörfer et al. (2002),

$$da_{i+\frac{1}{2}} = a_{i+1} - a_i,$$

is subject to large cancellation so that truncation errors inhibit the convergence of the Newton-Raphson scheme in the implicitly finite differenced time step. However, the problem can be avoided by a simple modification that was first explored in Fisker (2004). The state vector of Agile-Boltztran is given by the following set of quantities

$$y = (a, r, u, m, \rho, T, Y_e), \quad (4)$$

with enclosed baron mass a , radius r , velocity u , gravitational mass m , baryon density ρ , temperature T , electron fraction Y_e . In the improved version, the state vector at time t^n is based on zone masses, $da_{i+\frac{1}{2}}^n$, where the enclosed mass

$$a_i^n = \sum_1^{i-1} da_{i+\frac{1}{2}}^n$$

becomes the derived quantity.

The form of the generic equation (30) in Liebendörfer et al. (2002) applies to the continuity equation, the momentum equation and the energy equation. If we define $\delta_i = a_i^{n+1} - a_i^n$ as the difference of the enclosed mass a_i between time t^n and t^{n+1} , Eq. (30) in Liebendörfer et al. (2002) becomes

$$\frac{y_{i+\frac{1}{2}}^{n+1} \left(da_{i+\frac{1}{2}}^n + \delta_{i+1} - \delta_i \right) - y_{i+\frac{1}{2}}^n da_{i+\frac{1}{2}}^n}{dt} = y_{i+1}^{\text{adv}} - y_i^{\text{adv}} - y_{i+\frac{1}{2}}^{\text{ext}} = 0, \quad (5)$$

where the relative velocity between the fluid and the grid in the advection term y^{adv} is defined by

$$u_i^{\text{rel}} = -\frac{a_i^{n+1} - a_i^n}{dt} = -\frac{\delta_i}{dt}. \quad (6)$$

The cancellation of large numbers during the Newton-Raphson iterations of the implicit time step is avoided if the time shifts δ_i are chosen as the unknowns in the state vector when Eqs. (5) and (6) are solved. The vector of zone masses is then updated between the implicit time steps by

$$da_{i+\frac{1}{2}}^{n+1} = da_{i+\frac{1}{2}}^n + \delta_{i+1} - \delta_i.$$

This leads to satisfactory convergence of the Newton-Raphson iterations even in the presence of large density contrasts.

2.2 The equation of state

For the present investigation of the neutrino driven wind, the baryon EoS from Shen et al. (1998) for hot and dense nuclear matter has been implemented for matter in nuclear statistical equilibrium (NSE). For temperatures below $T = 0.5$ MeV where NSE does not apply, the baryon EoS combines an ideal gas approximation for a distribution of nuclei, based on Timmes and Arnett (1999) (including ion-ion-correlations), and a nuclear reaction network using the composition given by the progenitor model. Details of the reaction network are given in Thielemann et al. (2004) and references therein. The nuclear abundances are included in the state vector of Agile-Boltztran as follows

$$y = (a, r, u, m, \rho, T, Y_e, Y_1, \dots, Y_N). \quad (7)$$

For all these quantities, including the nuclear abundances $Y_1, \dots, Y_N$, the corresponding advection equations are solved as described in Liebendörfer et al. (2002) §3, but with an improved second order accurate total variation diminishing advection scheme. The nuclear reaction network is used in an operator-split manner in order to calculate the abundance changes due to the source terms which in turn depend on employed reaction rates.

Due to computational limitations, we restrict ourselves to $N = 19$. We consider the free nucleons and the 14 symmetric nuclei, from ^4He to ^{56}Ni . In order to model matter with $Y_e \neq 0.5$ to some extent, we additionally include ^{53}Fe , ^{54}Fe and ^{56}Fe . The network calculates the composition dynamically from the progenitor stage until the simulations are stopped. It is used for an accurate internal energy evolution. In addition, we can approximately reflect the composition of the PNS surface for more than 20 seconds after bounce, where nuclei that have been previously in NSE are freezing out of NSE as the temperature drops rapidly below 0.5 MeV already about 1 second after bounce and reach below 0.01

MeV at about 10 seconds post bounce. In previous studies the simplification of an ideal gas of Si-nuclei was used for matter which is not in NSE. This leads to an increasing inaccurate internal energy evolution after 500 ms post bounce when the explosion shock reaches the Si-layer of the progenitor and simplifications could not be extended beyond 1 second post bounce time. The implementation of the nuclear reaction network now makes it possible to include more mass (up to and including a large fraction of the He-layer, depending on the progenitor model) into the physical domain and follow the dynamical evolution of the explosion by one order of magnitude longer.

The baryon EoSs for NSE and for non-NSE are coupled to an electron-positron EoS (including photons), developed by Timmes and Arnett (1999).

2.3 Enhanced neutrino emissivity and opacity

By our choice of a spherically symmetric approach, we implement the explosion mechanism of massive Fe-core progenitor stars artificially to trigger neutrino driven explosions during the post bounce evolution after the deleptonization burst has been launched. Neutrino heating between the neutrinospheres and the SAS transfers energy from the radiation field into the fluid. A part of this energy is converted into thermal energy which revives the SAS and launches the explosion. The revival of the SAS and hence the neutrino driven explosions take place on a timescale of several 100 ms.

During the post bounce evolution, heavy nuclei continue to fall onto the SAS and dissociate into free nucleons. These free nucleons accrete onto the PNS surface. Hence the dominant neutrino heating contributions behind the SAS are due to the electronic charged current reactions Eqs. (1) and (2). To trigger explosions in spherically symmetric core collapse simulations of massive Fe-core progenitors, we enhance the emissivity j and absorptivity χ by a certain factor (typically 5 – 7) in the region between the SAS and the neutrinospheres. This corresponds to matter with entropies above 6 k_B /baryon and baryon densities below 10^{10} g/cm³. The entropies ahead of the shock are lower and the central densities of the PNS are higher, such that the artificial heating only applies to the region between the neutrinospheres and the SAS. The artificially enhanced reaction rates do not change the neutrino luminosities and mean neutrino energies significantly for the electron-neutrinos and electron-antineutrinos. Furthermore, β -equilibrium is fulfilled since the reverse reaction rates are obtained via the detailed balance. However, the timescale for weak-equilibrium to be established is reduced and hence the electron fraction changes on a shorter timescale. In combination with the increased neutrino energy deposition, this leads to a deviation of the thermodynamic variables in comparison to simulations using the standard opacities given in Bruenn (1985), which will be further discussed in §7. The weak neutrino driven explosions obtained have explosion energies of 6.5×10^{50} erg and 2×10^{50} erg for the 10.8 and the 18 $M_\odot$ progenitor model respectively.

2.4 Explosion energy and mass cut

The explosion energy estimate is a quantity which changes during the dynamical evolution of the system. It is given by the total specific energy of the fluid in the laboratory frame

$$E_{\text{specific}}(t, a) = \Gamma e + \frac{2}{\Gamma + 1} \left(\frac{u^2}{2} - \frac{m}{r} \right), \quad (8)$$

which in turn is the sum of the specific internal energy e ¹, the specific kinetic energy given by the fluid velocity $u = \partial r / \partial t$ squared and the specific gravitational binding energy m/r with gravitational mass m and radius r (see Liebendörfer et al. (2001b)). $\Gamma(t, a) = \sqrt{1 - 2m/r + u^2}$ and $\alpha(t, a)$ are the metric functions in a non-stationary and spherically symmetric spacetime with coordinate time t , baryon mass a and the two angular coordinates (θ, ϕ) describing a 2-sphere of radius $r(t, a)$ [see Misner and Sharp (1964)]. The explosion is determined by the energy of the ejecta. Integrating $E_{\text{specific}}(t, a)$ with respect to the enclosed baryon mass starting from the progenitor surface M toward the center

$$E_{\text{total}}(t_0, a_0) = - \int_M^{a_0} E_{\text{specific}}(t_0, a) da, \quad (9)$$

¹The baryon contribution to the internal energy is composed of a thermal and nuclear part, i.e. $e = e_{\text{thermal}} + e_{\text{nuclear}}$. In NSE, e is given implicitly via the EoS of hot and dense nuclear matter. In non-NSE, e_{nuclear} is the binding energy of the nuclear composition used in the reaction network.

gives the total mass-integrated energy, at a given time t_0 outside a given mass a_0 . The expression (9) is negative during the collapse, bounce and the early post bounce phases because the progenitor and central Fe-core are gravitationally bound. At some time after bounce, expression (9) becomes positive in the region between the shock and the neutrinospheres. It stays negative at large distances and close to the deep gravitational potential of the PNS, because the outer layers of the progenitor and the PNS continue to be gravitationally bound. While the emission of neutrinos cools the PNS interior, neutrino absorption deposits energy on a timescale τ_{heating} of the order 100 ms into the fluid near the neutrinospheres. This increases the specific internal energy which matches at later (~ 500 ms) post bounce times the gravitational binding energy at a certain distance toward the center.

On a suggestion by S. Bruenn, we consider the *mass cut* as follows

$$a_{\text{cut}} = a \left(\max \left(E_{\text{total}}(t \gg \tau_{\text{heating}}, a) \right) \right). \quad (10)$$

The material outside of a_{cut} is gravitationally unbound and will be ejected while the enclosed material will accrete onto the central PNS. The explosion energy estimate is defined as the total mass-integrated energy of the layers outside the mass cut

$$E_{\text{expl}} = E_{\text{total}}(t \gg \tau_{\text{heating}}, a_{\text{cut}}), \quad (11)$$

at late times ($t \gg \tau_{\text{heating}}$) after the explosion has been launched. It becomes clear from the above expressions that the explosion energy estimate is sensitively determined by the balance of internal and kinetic energies to gravitational binding energy.

From the time post bounce when the shock reaches low enough densities and temperatures such that neutrinos decouple from matter completely, neutrino heating and cooling does not affect the explosion energy estimate anymore. The additional energy deposition from the neutrino driven wind, which will be discussed further below, might affect the explosion estimate at later times. We will illustrate in particular the effect of the formation of a supersonic neutrino driven wind and the wind termination shock to the explosion energy estimate. Only after the neutrino driven wind disappears, the final value of the explosion energy can be obtained.

2.5 The neutrino observables

The neutrino radiation hydrodynamics equations are a coupled system which combines the evolution of the matter properties given by the state vector y and the radiation field, as documented in Liebendörfer et al. (2004) and references therein. The neutrino radiation field is taken into account via the phase-space distribution function $f_\nu(t, a, \mu, E)$ for each neutrino flavor $\nu = (\nu_e, \bar{\nu}_e, \nu_{\mu/\tau}, \bar{\nu}_{\mu/\tau})$. In spherical symmetry, it depends on the time t , the enclosed baryon mass a as well as on the neutrino energy E and the cosine of the propagation angle $\mu = \cos(\theta)$. The evolution of the distribution function is modeled via solving the Boltztran transport equation. It determines the phase-space derivative of the specific distribution function $F_\nu = f_\nu/\rho$, i.e. the distribution function divided by the matter density ρ , in a co-moving frame (see for example Eq.(8) in Liebendörfer et al. (2005)) and due to neutrino-matter interactions such as emission and absorption as well as scattering and pair reactions.

In order to compare simulation results, neutrino observables can be defined. Commonly used is the neutrino number-luminosities, which is given by the phase-space integration of the neutrino distribution function as follows

$$L_n([t_0, t_1], a) = 4\pi r^2 \rho \frac{2\pi c}{(hc)^3} \int_{-1}^{+1} d\mu \int_0^\infty E^2 dE F_\nu(t, a, \mu, E),$$

which is the number of neutrinos ² of energy E passing through the mass coordinate a for a given time-interval $[t_0, t_1]$ taken in a co-moving frame at position $r(t, a)$. Equivalently, the energy-luminosity can be defined as follows

$$\begin{aligned} L_e([t_0, t_1], a) &\equiv L_\nu([t_0, t_1], a) \\ &= 4\pi r^2 \rho \frac{2\pi c}{(hc)^3} \int_{-1}^{+1} \mu d\mu \int_0^\infty E^3 dE F_\nu(t, a, \mu, E), \end{aligned}$$

²The integration with respect to μ is performed separately for in- and out-ward direction, according to the transport coefficients.

for each neutrino flavor, i.e. $(\nu_e, \bar{\nu}_e, \nu_{\mu/\tau}, \bar{\nu}_{\mu/\tau})$. Additionally useful quantities are the mean neutrino and root-mean-squared (rms) neutrino energies, defined as follows

$$\begin{aligned}\langle E_\nu(t, a) \rangle &= \frac{\int_{-1}^{+1} d\mu \int_0^\infty E^3 dE F_\nu(t, a, \mu, E)}{\int_{-1}^{+1} d\mu \int_0^\infty E^2 dE F_\nu(t, a, \mu, E)}, \\ \langle E_\nu(t, a) \rangle_{\text{rms}} &= \sqrt{\frac{\int_{-1}^{+1} d\mu \int_0^\infty E^4 dE F_\nu(t, a, \mu, E)}{\int_{-1}^{+1} d\mu \int_0^\infty E^2 dE F_\nu(t, a, \mu, E)}}.\end{aligned}$$

We will use these observables, i.e. the energy-luminosities and the mean and root-mean-squared energies, to illustrate the dynamical evolution of the radiation field as well as for comparisons with previous studies.

2.6 The electron fraction

The proton-to-baryon ratio is essential for the composition of the ejecta, which is obtained via detailed post processing nucleosynthesis calculations. In the absence of muons or taus, the proton-to-baryon ratio is given by the electron fraction as follows

$$Y_e = Y_{e^-} - Y_{e^+}, \quad (12)$$

which defines the total number of charges per baryon. The change of the electron fraction is given by the balance of emitted and absorbed electrons (positrons) and electron-neutrinos (antineutrinos) at free nucleons and nuclei. Weak-equilibrium is achieved if

$$\mu_{e^-} + \mu_p = \mu_n + \mu_{\nu_e}, \quad (13)$$

$$\mu_{e^+} + \mu_n = \mu_p + \mu_{\bar{\nu}_e}, \quad (14)$$

where μ_i are the chemical potentials for electron and positron ($\mu_{e^\pm}$), proton (μ_p), neutron (μ_n) and electron-neutrino (μ_{ν_e}) and electron-antineutrino ($\mu_{\bar{\nu}_e}$). The time-derivative of the electron fraction, $\dot{Y}_e$, is given by the phase-space integration of the emissivities j_ν and the absorptivities χ_ν for electron-neutrinos and electron-antineutrinos as follows

$$\begin{aligned}\dot{Y}_e &= -\frac{2\pi}{(hc)^3} \frac{m_B c}{\rho} \int_{-1}^{+1} d\mu \int_0^\infty E^2 dE \times \\ &\times ((j_{\nu_e} - \tilde{\chi}_{\nu_e} f_{\nu_e}) - (j_{\bar{\nu}_e} - \tilde{\chi}_{\bar{\nu}_e} f_{\bar{\nu}_e}))\end{aligned} \quad (15)$$

where m_B is the baryon mass, ρ is the matter density and $\tilde{\chi} = j + \chi$. The emissivities $j_\nu(E)$ and absorptivities $\chi_\nu(E)$ depend on the neutrino energy. They are the reaction rates for the electronic charged current reactions, which are calculated following Bruenn (1985) and depend on temperature and density. Eq. (15) is found by combining the equation of lepton number conservation with the phase-space integrated Boltzmann transport equation (see Mezzacappa & Bruenn (1993 a)). To follow the dynamical evolution of the electron fraction via Eq. (15), knowledge of the neutrino distribution functions is required for which neutrino transport is necessary.

3 Explosions in spherical symmetry

Progenitor stars more massive than $9 M_\odot$ develop extended Fe-cores at the end of stellar evolution. The explosion mechanism of such Fe-core progenitors is an active subject of research. In the following section, we will investigate the neutrino driven explosions of the 10.8 and $18 M_\odot$ Fe-core progenitors from Woosley et al. (2002) in spherical symmetry by enhancing the electronic charged current reaction rates artificially. Further below, we will investigate the explosion phase of the $8.8 M_\odot$ O-Ne-Mg-core from Nomoto (1983, 1984, 1987), where the explosion is obtained using the standard neutrino opacities as introduced in § 2.

3.1 Neutrino driven explosions of Fe-core progenitors

The dynamical behavior of the explosion energy estimate and the shock position are illustrated in Figs. 1 (a) and (b) respectively with respect to time after bounce. The figures illustrate the explosion phase and long time evolution up to 10 seconds after bounce. After achieving a convergent value between 600 ms and 2 seconds post bounce of 4.5×10^{50} erg, the explosion energy estimate is lifted slightly to about 6.5×10^{50} erg. This effect coincides with the additional mass outflow obtained in the neutrino driven wind and the appearance of the reverse shock, which will be discussed further below. In simulations with a less intense (subsonic) neutrino driven wind, this effect is much weaker and the explosion energy can be obtained already after about 1 second post bounce.




Figure 1: Explosion energy estimate and shock position with respect to time after bounce for the $10.8 M_{\odot}$ progenitor model from Woosley et al. (2002). In addition, graph (b) illustrates the position of the He-layer (dashed line) and the O-layer (dash-dotted line).

The neutrino luminosities and the mean as well as rms neutrino energies are shown in Fig.2 for the 10.8 (middle panel) and the 18 (right panel) $M_{\odot}$ progenitor model with respect to time after bounce. Note that the more compact PNS of the $18 M_{\odot}$ progenitor model results in generally larger neutrino luminosities. The oscillating shock position and the consequent contracting and expanding neutrinospheres during the neutrino heating phase of the 10.8 and $18 M_{\odot}$ progenitor models on a timescale of several 100 ms are reflected in the electron-flavor neutrino luminosities, which correspondingly increase and decrease respectively. During the heating phase, the mean neutrino energies of the electron-(anti)neutrinos increase from about 8 (10) MeV after bounce to about 12 (14) MeV until the explosion is launched. The mean neutrino energy of the μ/τ -neutrinos remains constant at about 18 MeV during the heating phase. The mean neutrino energies are generally smaller than the rms-energies but follow the same behavior. The explosions for both models are launched after about 350 ms post bounce, which defines the neutrino heating timescale for the energy deposition in the gain region to revive the SAS. Matter is accelerated to positive velocities and the SAS turns into the dynamic explosion shock. The resulting neutrino spectra from these artificially induced explosions in spherical symmetry are in general agreement with the neutrino spectra from axially-symmetric neutrino driven core collapse supernova models that explode without artificially modified reaction rates (see Marek and Janka (2009)). The explosion shock continuously propagates through the remaining domain of the progenitor star. After the explosions have been launched, the electron flavor neutrino luminosities decay exponentially. Furthermore the jumps in the neutrino energies

after 350 ms post bounce for the 10.8 and 18 $M_{\odot}$ progenitor models are due to the shock propagation over the position of 500 km, where the observables are measured in a co-moving reference frame.




Figure 2: Neutrino luminosities and energies with respect to time after bounce for the 8.8 $M_{\odot}$ O-Ne-Mg-core progenitor model from Nomoto (1983,1984,1987) (left panel) and the 10.8 $M_{\odot}$ and 18 $M_{\odot}$ Fe-core progenitor models from Woosley et al. (2002) (middle and right panels respectively), measured in a co-moving frame at 500 km distance.

3.2 The O-Ne-Mg-core

A special star is the 8.8 $M_{\odot}$ progenitor model from Nomoto (1983,1984,1987). The central thermodynamic conditions at the end of stellar evolution are such that only a tiny fraction of about 0.15 $M_{\odot}$ of Fe-group nuclei are produced, where nuclear statistical equilibrium (NSE) applies (see Fig. 3 (a) top panel). Instead, the central composition is dominated by ^{16}O , ^{20}Ne and ^{24}Mg nuclei. Because temperature and density increase during the collapse, these nuclei are burned into Fe-group nuclei and the NSE regime increases (see Fig. 3 middle panel). The core continues to deleptonize, which can be identified at the decreasing Y_e in Fig. 3. We use our nuclear reaction network as described in §2.2 to calculate the dynamically changing composition, based on the abundances provided by the progenitor model. The size of the

bouncing core of $M_{\text{core}} \approx 0.65 M_{\odot}$ is significantly larger in comparison with the previous studies by Kitaura et al. (2006) and Liebendörfer (2004), illustrated in Fig. 4(a) at different velocity profiles before and at bounce. This is because we do not take the improved electron capture rates from Hix et al. (2003) and Langanke et al. (2003) into account, which are based on the capture of electrons on the distribution of heavy nuclei. It results in a lower central electron fraction at bounce and a consequently more compact bouncing core of about $\approx 0.1 M_{\odot}$, in comparison to the standard rates given in Bruenn (1985). The remaining difference is most likely due to the large asymmetry energy of the EoS from Shen et al. (1998) applied to the present study.

This progenitor is not only a special case due to the incomplete nuclear burning at the end of stellar evolution but also due to the steep density gradient which separates the dense core from the He- and H-envelope at $1.376 M_{\odot}$, see Fig. 3 (c). There, the density drops over 13 orders of magnitude which makes it difficult to handle numerically.




Figure 3: Evolution of the $8.8 M_{\odot}$ progenitor model from Nomoto (1983,1984,1987) during the core collapse phase (top: progenitor configuration, middle: 24 ms before bounce, bottom: at bounce). The composition (left panels) are as follows: C+O (thin solid line), Ne+Mg (dashed line), Fe+Ni (dash-dotted line), He (dotted line), Si+S (thick solid line).

The low density of the mass outside the O-Ne-Mg-core makes it possible to obtain the explosion in spherical symmetry supported via neutrino heating. Neutrino cooling in the region of dissociated nuclear matter causes the expanding shock front to turn into the SAS with no significant matter outflow. ν_e -cooling dominates over $\bar{\nu}_e$ -heating by one order of magnitude. Only at the dissociation line of infalling heavy nuclei, the neutrino energy deposition drives the SAS slowly to larger radii, for illustration see the heating(cooling) rates and velocity profile in Fig. 5 (left panel) at




Figure 4: Radial velocity profiles with respect to the baryon mass (a) and with respect to the radius (b) for the $8.8 M_{\odot}$ progenitor model from Nomoto (1983,1984,1987)

20 ms post bounce. However, the cooling of ν_e still contributes to a large amount at 25 ms post bounce over the heating of $\bar{\nu}_e$ and $\nu_{\mu/\tau}$ in Fig. 5 (middle panel) behind the SAS. Only directly at the shock a small net-heating rate remains. Hence the influence of the neutrinos to the explosion itself is of minor importance. More important is the region of C-O-burning which produces Ne and Mg. The hydrodynamic feedback to this thermodynamic transition can be identified already during the collapse phase of the progenitor core at the velocity profiles in Fig. 4(a) at about $1.35 - 1.374 M_{\odot}$. As material is shock heated post bounce, the transition layer where Ne and Mg nuclei are burned into NSE propagates together with the expanding shock wave outwards. In other words, the Ne-Mg-layer of the progenitor is converted directly into NSE. Furthermore, the transition (discontinuity) from C-O-burning is falling quickly towards the SAS. It was found to be at about 350 km at 20 ms post bounce and at about 200 km at 25 ms post bounce, illustrated at the velocity profiles (bottom) in Fig. 5 (left-right panels). At about 30 ms post bounce, the entire Ne-Mg-layer is converted into NSE due to the temperature increase obtained via shock heating. Hence, C and O nuclei are burned directly into NSE.

In contrast to more massive Fe-core progenitors where O-burning produced an extended Si-S-layer, the amount of ^{28}Si and ^{32}S is less than 1% at the end of nuclear burning for the O-Ne-Mg-core discussed here (see Fig. 3 (a) middle panel). This small fraction of Si and S is already converted into NSE during the initial collapse phase, due to the rapid density and temperature increase of the contracting core. Hence, C- and O-nuclei are burned directly into NSE during the post bounce evolution. This sharp transition is related to a jump in the density and the thermodynamic variables. As the SAS propagates over this transition along the decreasing density, the shock accelerates to positive velocities (see Fig. 5 right panel). The consequent explosion is hence driven due to the shock propagation over the infalling transition between two different thermonuclear regimes rather than by pure neutrino heating, illustrated at the velocity profiles in Fig. 4(b). Although Kitaura et al. (2006) approximated nuclear reactions during the evolution of the O-Ne-Mg-core progenitor, the results of their explosion dynamics are in qualitative agreement with our findings.

The more massive Fe-core progenitors show the same thermo- and hydrodynamic features as discussed above for the O-Ne-Mg-core due to the transition between different thermonuclear regimes. However, the differences are smaller because C-O-burning produces an extended layer composed of ^{28}Si and ^{32}S . The transition of Si-burning into NSE is much smoother than the transition of C-O-burning into NSE. Furthermore, due to the more massive Si-S and C-O-layers for the Fe-core progenitors, the transitions take more time of the order 100 ms to fall onto the SAS. The presence of neutrino heating becomes important for the more massive Fe-core progenitors to drive the SAS to large radii on a longer timescale. The effects of the shock propagation across the transition between different thermonuclear regimes has been pointed out in Bruenn et al. (2006) with respect to the explosion dynamics in axially-symmetric simulations of massive Fe-core progenitors. In our spherically symmetric model we can not confirm the driving force of explosions of Fe-core progenitors to be the shock propagation across different thermonuclear regimes. It is rather the neutrino energy deposition, although enhanced, which initiates the explosion. The shock is accelerated additionally when crossing different thermonuclear regimes due to the density jumps at the transitions.




Figure 5: Heating (> 0) and cooling (< 0) rates for the $8.8 M_{\odot}$ progenitor model from Nomoto (1983,1984,1987) during the explosion phase at 20 ms (left panel), 25 ms (middle panel) and 30 ms (right panel) after bounce. For a better comparison, the velocities are plotted for the same post bounce times.

3.3 Comparison of the neutrino spectra

Striking is the agreement in the mean neutrino energies between all different progenitor models (including the O-Ne-Mg-core and the Fe-core progenitors) during the explosion phase, although the neutrino emissivities and opacities are enhanced for the Fe-core progenitor models (see Fig. 2). The explosion phase for the O-Ne-Mg-core lasts only until about 40 ms post bounce and is significantly shorter in comparison to the more massive Fe-core progenitors. Furthermore, the luminosities are also smaller by a factor of 2. For all models, the electron antineutrino luminosity is larger than the electron neutrino luminosity on a timescale of 200 ms after the explosion has been launched. This slight difference reduces again at later times where the electron neutrino luminosity becomes again larger than the electron antineutrino luminosity. However, after the explosion has been launched the behavior of the luminosities are in qualitative agreement for all models. The same holds for the mean neutrino energies which increase continuously during the neutrino heating phase. The electron (anti)neutrinos have rms energies of about 12 (14) MeV where as after the explosion has been launched, rms energies of about 11 (13) MeV are obtained. The values remain constant on the timescale of 1 second post bounce. The (μ/τ) -neutrinos have rms energies of about 18 MeV during the neutrino heating phase and about 15 MeV after the explosion has been launched. These differences in the mean neutrino energies and luminosities during the neutrino heating, initial and proceeding explosion phases are in correspondence with the electron fraction of the material, as will be illustrated in the following section.

3.4 The electron fraction of the early ejecta

During the neutrino heating phase, the neutrino spectra are mainly determined by mass accretion at the neutrinospheres. Neutron rich nuclei from the progenitor star with an electron fraction of $Y_e \approx 0.45$ are falling onto the oscillating SAS and dissociate into free nucleons and light nuclei, see Fig. 6 (d). These nucleons accrete than slowly onto the PNS surface at the center. Due to the increased electron-degeneracy behind the SAS in Fig. 6 (c), weak-equilibrium is

established at a lower value of the electron fraction of $Y_e \leq 0.15$.




Figure 6: Selected hydrodynamic variables during the initial explosion phase at three different post bounce times for the $10.8 M_{\odot}$ progenitor model from Woosley et al. (2002).

As soon as the SAS is revived and propagates outward, see the velocity and density profiles in Fig. 6 (a) and (b), the electron degeneracy behind the expanding shock is reduced and weak-equilibrium is established at a larger value of the electron fraction of $Y_e > 0.56$ ³. The capture rates for electron-neutrinos at neutrons are favored over electron-antineutrino captures at protons. This slight difference results in an electron and hence proton excess. Consequently the explosion ejecta are found to be initially proton-rich. This behavior of the electron fraction was found for all our explosion models, for the 10.8 and $18 M_{\odot}$ Fe-core progenitors with artificially enhanced opacities and for the O-Ne-Mg-core using the standard opacities. Such explosion models were investigated with respect to the nucleosynthesis in general and with respect to the νp -process by Fröhlich et al. (2006a-c).

One of the main goals of the present investigation is to determine the behavior of the electron fraction for the initially proton-rich ejecta on a long timescale of the order 10 seconds, in a consistent manner. We explore the question if the material ejected in the neutrino driven wind becomes neutron-rich and which are the conditions (e.g. entropy per baryon, and expansion timescale) obtained in the neutrino driven wind. These aspects are of special relevance for the composition of the ejecta, which is determined via explosive nucleosynthesis analysis, in particular in order to be able to draw conclusions with respect to a possible site for the production of heavy elements via the r -process. Therefore, the continued expansion of the explosion ejecta must be simulated, for which the inclusion of a large physical domain of the progenitor up to the He-layer is required. Furthermore, since only the electronic charged current reactions including the neutrino fluxes can determine the electron fraction, the PNS contraction at the center and hence the contraction of

³ The EoS from Shen et al. (1998) is limited to a maximum electron fraction of $Y_e \geq 0.564$. The EoS has been extended by Gögelein (2007) to model asymmetric nuclear matter with an electron fraction above 0.564. We assume an ideal gas of free nucleons and light nuclei, which is a sufficient assumption for the conditions found in the region of the extremely proton-rich ejecta.

the neutrinospheres are essential.

4 The neutrino driven wind

In this section we investigate the post explosion evolution with special focus on the properties of the ejecta, in particular the electron fraction. We explore the problem if the initially proton-rich ejecta become neutron rich at later times of the order 10 seconds and if the conditions might indicate a possible site for the nucleosynthesis of heavy nuclei via the *r*-process. This has been assumed in static steady-state as well as parametrized dynamic wind models, based on the results obtained by Woosley et al. (1994). The ejected material in their simulations does never become proton-rich, the electron fraction was found to continuously decrease with time after the explosion has been launched. Thus, the ejecta were investigated in a region where the conditions are favorable for the *r*-process. The stellar models applied to the present investigation of the neutrino driven wind are the 8.8 $M_{\odot}$ O-Ne-Mg-core and the 10.8 and 18 $M_{\odot}$ Fe-core progenitors, where for the latter two cases the explosions are obtained using the artificially enhanced opacities as described in §2.3.

After the explosion has been launched, the region between the expanding explosion shock and the neutrinospheres cools rapidly and the density decreases continuously as illustrated in Figs. 7 and 8 (d) and (h). In order to determine the evolution of the electron fraction Y_e , the non-local neutrino fluxes are as important as the local neutrino reaction rates. Since the PNS and hence the neutrinospheres contract continuously due to the deleptonization, the degeneracy increases and matter at the PNS surface is found to be extremely neutron-rich with $Y_e \leq 0.1$ (see Figs. 7 and 8) (d) and (f).

Independent of the progenitor model, the region on top of the PNS surface is continuously subject to neutrino heating during the post explosion phase. The dominant heating sources are the absorption of electron-(anti)neutrinos on free nucleons, due to the large fraction of free nucleons (dissociated nuclear matter) in the region on top of the PNS, as shown in Fig. 9. The neutrino pair production and thermalization processes have a negligible contribution to the heating outside the neutrinospheres. In order to compare the heating and cooling rates in Fig. 9, we plot the quantities with respect to the baryon density. While neutrino cooling is still dominantly present at ~ 500 ms post bounce (thin lines in Fig. 9), at later times after ~ 1 second post bounce (thick lines in Fig. 9) neutrino cooling vanishes and only heating is found in the density domain of interest, i.e. between $10^7 - 10^{12}$ g/cm³. Figs. 7 (d) and 8 (d) show the conditions for the contracting PNSs at the center via the radial baryon density profiles and the electron-neutrinospheres. The region of interest where the neutrino driven wind develops corresponds to the density domain of $10^7 - 10^{10}$ g/cm³. The degeneracy of the early ejecta favors proton-rich matter where a large electron fraction of $Y_e \approx 0.54$ is obtained. Hence, the absorption of electron-antineutrinos on free protons dominates over electron-neutrino absorption on free neutrons. The corresponding radial neutrino luminosities and rms energies are shown in Figs. 7 and 8 (e) and (g). In addition, for the first time we were able to follow the deleptonization burst from core bounce for several seconds over a large physical domain including several 10^5 km of the progenitor star. The outrunning luminosity peak can be identified in the luminosities in Figs. 7 and 8 (e) at 0.6 seconds after bounce at a distance between 5×10^4 and 10^5 km, leaving the computational domain between 1 – 2 seconds post bounce.

After the explosion has been launched, the continued energy transfer from the neutrino radiation field into the fluid outside the neutrinospheres as illustrated in Fig. 9 drives the matter entropies to large values, shown in Figs. 7 and 8 (c). The heat deposition at the PNS surface accelerates matter to positive velocities, see Figs. 7 and 8 (a), after ≈ 1 second post bounce. This matter outflow is known as the neutrino driven wind, which proceeds adiabatically at larger radii. This is consistent with the constant radial entropy per baryon profiles in the graphs (c). Furthermore, the rapidly decreasing luminosities reach values below 5×10^{51} erg/s already 1 second after bounce (see Figs. 7 and 8 (e)). The luminosities continue to decrease and reach values below 1×10^{51} erg/s at 10 seconds after bounce. The mean neutrino energies also decrease constantly where values below 10 MeV for the electron-flavor neutrinos and below 12 MeV for the (μ/τ) -neutrinos are obtained (see Figs. 7 and 8 (g)).

Several previous wind studies achieved supersonic matter outflow velocities for the neutrino driven wind due to assumed large luminosities. In any case, the accelerated material of the neutrino driven wind collides with the slower and subsonically expanding explosion ejecta. Comparing Figs. 7 and 8, the more compact wind region of the 18 $M_{\odot}$ progenitor model produces a less pronounced neutrino driven wind in comparison to the 10.8 $M_{\odot}$ progenitor model. The densities of the wind region are larger up to two orders of magnitude and the temperatures are higher by a factor




Figure 7: Selected hydrodynamic variables during the formation of the neutrino driven wind at three different post bounce times for the $10.8 M_{\odot}$ progenitor model from Woosley et al. (2002). In addition, graphs (e) and (g) show the neutrino luminosities and rms neutrino energies (solid lines: ν_e , dashed lines: $\bar{\nu}_e$, dash-dotted lines: $\nu_{\mu/\tau}$). For this progenitor model the neutrino driven wind becomes supersonic, using the enhanced opacities.




Figure 8: The same configuration as Fig. 7 for the $18 M_{\odot}$ progenitor model from Woosley et al. (2002). The neutrino driven wind remains subsonic for this progenitor model, even using the enhanced opacities.




Figure 9: Heating and cooling rates with respect to the baryon density at three different post bounce times 0.5 second (thin lines), 1 second (intermediate lines), 5 seconds (thick lines) for the $10.8 M_{\odot}$ progenitor model from Woosley et al. (2002).

of 2. The resulting velocities of the neutrino driven wind outflow are smaller by a factor of 2 and stay below 10^4 km/s. Hence, the neutrino driven wind remains subsonic for all times for the $18 M_{\odot}$ progenitor model (see Fig. 8 (a)) where the wind develops supersonic velocities for the $10.8 M_{\odot}$ progenitor model (see Fig. 7 (a)). In the case of a supersonic neutrino driven wind, this leads to the formation of a reverse shock known as the wind termination shock. The formation of the wind termination shock of the $10.8 M_{\odot}$ progenitor model is illustrated in Fig. 10 and will be discussed in the following paragraph. In order to analyze the dynamical evolution and the consequences of the formation of the reverse shock, steady-state wind models can not be used. Radiation hydrodynamics is required in order to describe the dynamical effects consistently. Our results, obtained using general relativistic radiation hydrodynamics based on spectral three flavor Boltzmann neutrino transport, are in qualitative agreement with the detailed parametrized investigation by Arcones et al. (2007).

During the initial and subsonic wind expansion, the matter entropies in Fig. 10 (c) increase slowly from 4 to 5 – 10 k_B /baryon and the densities in Fig. 10 (b) and temperatures in Fig. 10 (e) decrease on a long timescale over several seconds. Furthermore, the reduced degeneracy in the wind increases the electron fraction shown in Fig. 10 (d) slowly on the same timescale. When the material is accelerated supersonically with velocities of several 10^4 km/s up to radii of a several 10^3 km (see Figs. 10 (a) and (f)), the entropies increase from $s \approx 5 - 10 k_B$ /baryon to $s \approx 40 - 60 k_B$ /baryon on a short timescale of the order of 100 ms. During this rapid expansion, the density and temperature decrease drastically from 10^{10} g/cm³ to $10^4 - 10^2$ g/cm³ and from 3 MeV to 0.1 – 0.01 MeV respectively (see Figs. 10 (b) and (e)). It also corresponds to a rapid decrease of the degeneracy which in turn is reflected in a rapid increase of the electron fraction of the accelerated material on top of the PNS surface, from $Y_e \approx 0.1$ to $Y_e \approx 0.56$ (see Fig. 10 (d)). Furthermore, the supersonically expanding neutrino driven wind collides with the explosion ejecta as can be seen in Fig. 10 (a) (solid red line) at radii of several 10^4 km. Consequently, the previously accelerated material decelerates behind the explosion ejecta as can be seen in the velocities in Fig. 10 (f). This phenomenon becomes significant after about 2 seconds post bounce and corresponds to the formation of the reverse shock, i.e. the wind termination shock. (see Fig. 10 (a) dashed red line at radii of several 10^3 km). It causes an additional entropy increase to the final values of $s \approx 50 - 100 k_B$ /baryon. During the rapid deceleration on the same short timescale of the order 100 ms, the densities in Fig. 10 (b) and temperatures in Fig. 10 (e) increase again slightly, where the degeneracy increases and hence the electron fraction reduces slightly to values of $Y_e \approx 0.54$. The following dynamical evolution is given by the subsonic and adiabatic expansion of the explosion ejecta on a longer timescale of the order of seconds. The density and temperature decrease slowly where the entropies of $s \approx 50 - 100 k_B$ /baryon and the electron fraction of about $Y_e = 0.54$ remain constant. The latter aspects are essential for the nucleosynthesis analysis of the ejecta. It can be understood in the sense that the neutrino reaction rates freeze out and the matter conditions correspond to the neutrino free streaming regime.

Note that the strong neutrino driven wind for the $10.8 M_{\odot}$ progenitor model is obtained using the enhanced opacities as introduced in §2.3. We additionally illustrate selected properties of the neutrino driven wind for the $8.8 M_{\odot}$ progen-

itor model in Fig. 11 where a strong neutrino driven wind was obtained using the standard emissivities and opacities given in Bruenn (1985). This is due to the low density of the region between the neutrinospheres at the PNS surface and the expanding explosion shock, where neutrino heating via the standard rates and energy from nuclear burning are sufficient to drive a strong supersonic matter outflow. Matter entropies increase to $s \approx 10 \text{ k}_B/\text{baryon}$ during the initial acceleration of the wind and the densities and temperatures decrease slowly on a timescale of seconds. The properties during the initial acceleration observed are similar to those of the more massive $10.8 M_\odot$ Fe-core progenitor. The same holds for the acceleration to supersonic velocities. The timescale is reduced to 100 ms where the entropies increase rapidly to $s \approx 20 - 50 \text{ k}_B/\text{baryon}$ (see Fig. 11 (c)) and due to the reduced degeneracy the electron fraction increases from $Y_e = 0.1$ at the PNS surface to $Y_e = 0.56$ (see Fig. 11 (d)). Density and temperature decrease to $10 - 100 \text{ g/cm}^3$ and 0.001 MeV respectively (see Fig. 11 (b) and (e)). The difference to the more massive $10.8 M_\odot$ Fe-core progenitor is due to the smaller mass enclosed between the PNS surface and the expanding explosion ejecta. For the more massive $10.8 M_\odot$ Fe-core progenitor in Fig. 10 (f), the previously accelerated material collides with the explosion ejecta already after a few 100 ms. Here the supersonic wind expands on a much longer timescale up to several seconds before it collides with the explosion ejecta (see Fig. 11 (f)). During this adiabatic expansion, entropy and electron fraction remain constant. The fast material collides with the much slower expanding explosion ejecta so that the material is decelerated and the reverse shock appears. This is again similar to the formation of the reverse shock for the more massive $10.8 M_\odot$ Fe-core progenitor as discussed above. Matter entropies increase to several $100 \text{ k}_B/\text{baryon}$ (see Fig. 11 (c)), density and temperature increase slightly (see Fig. 11 (b) and (e)) and the electron fraction reduces slightly to $Y_e \approx 0.52 - 0.54$ due to the increased degeneracy (see Fig. 11 (d)). The following evolution is determined by the adiabatic expansion of the explosion ejecta during which the entropy and electron fraction remain constant.

In the following paragraph, we will discuss the composition of the neutrino driven wind region to some extent. This is possible due to the recently implemented nuclear reaction network. It includes the free nucleons and the symmetric nuclei from ^4He to ^{56}Ni plus ^{53}Fe , ^{54}Fe and ^{56}Fe . The initial composition is given by the progenitor model. Mostly ^{28}Si and ^{30}S are shock-heated and burned to Fe-group nuclei due to the temperature and density jump during the initial expansion of the explosion shock (see Fig. 12 and compare with Figs. 7 and 8 (d) and (f)). The large fraction of these Fe-group nuclei reduces behind the explosion shock due to photodisintegration, indicated by the region of low density and high entropy in Figs. 7 and 8 (b) and (c). This produces a large fraction of α -particles, which in our model represent light nuclei. The region of α -particle domination behind the expanding explosion shock increases with time. This behaviour is illustrated in Fig. 12 for both Fe-core progenitor models under investigation. The position of the explosion shock coincides with the maximum of the mass fraction of Fe-group nuclei (in particular ^{56}Ni). In addition, density and temperature of the neutrino driven wind on top of the PNS surface decrease continuously with time. The low temperatures and densities in that region do not justify the assumption of NSE beyond ~ 1 second after bounce, where temperatures reach values below 0.5 MeV . Instead, our nuclear reaction network is used to determine the composition in that region. The decreasing density and temperature and the presence of a large fraction of free nucleons favor the freeze out of light nuclei. Finally, the entire region between the expanding explosion shock and the PNS surface is found to be dominated in our simulations by α -particles. In Fig. 12, the radii of the NSE to non-NSE transitions are indicated by vertical lines. The slight mismatch between the abundances between the heavy 'Fe'-group nuclei (the representative heavy nucleus with average atomic mass and charge in NSE) and ^{56}Ni (non-NSE) as well as between the α 's is due to the different nuclear models used for the two regimes. While in NSE the EoS for hot and dense nuclear matter assumes ^{56}Fe as the most stable nucleus due to the smallest mass per nucleon for low temperatures and densities, the nuclear reaction network applied in non-NSE calculates the composition dynamically based on tabulated reaction rates.

5 Comparison with previous wind studies

5.1 The proton-to-baryon ratio of the wind

The approximations made in previous wind studies is a simplified description of the radiation-hydrodynamics equations, see for example Duncan et al. (1986) and Qian and Woosley (1996). More crucial is the absence of neutrino transport. Neutrino heating and cooling is calculated only based on parametrized neutrino luminosities and mean energies. Hence, such models explore the neutrino driven wind by varying the neutrino luminosities and energies, where




Figure 10: Evolution of selected mass shells in the neutrino driven wind (as listed in graph (a) from 1.4429 – 1.4437 $M_{\odot}$ baryon mass) for the $10.8 M_{\odot}$ progenitor model from Woosley et al. (2002) where the enhanced opacities are used. Graph (a) shows in addition gray-scaled the entropy per baryon, the position of the expanding explosion shock (thick solid line) and the position of the wind termination shock (thick dashed line).




Figure 11: Evolution of selected mass shells in the neutrino driven wind (as listed in graph (a) from 1.34718 – 1.34750 $M_{\odot}$ baryon mass) for the 8.8 $M_{\odot}$ progenitor model from Nomoto (1983,1984,1987) where the standard emissivities and opacities given in Bruenn (1985) are used. The graphs show the same configurations as Fig. 10.




Figure 12: Selected radial mass fraction profiles during the neutrino driven wind phase for the $10.8 M_{\odot}$ (left panel) and the $18 M_{\odot}$ (right panel) progenitor model from Woosley et al. (2002) at 1 second (top), 3 seconds (middle) and 5 seconds (bottom) post bounce. The vertical lines represent the separation of NSE (EoS for hot and dense nuclear matter) where heavy nuclei are represented by a single Fe-group nucleus 'Fe' with average atomic mass and charge and non-NSE (nuclear reaction network) where the most abundant Fe-group element is ^{56}Ni , at temperatures of ≈ 0.5 MeV.

the simplified radiation-hydrodynamics equations are solved (see for example Thompson et al. (2001)). Since neutrino transport is neglected, the evolution equation for the electron fraction Eq. (15) cannot be solved consistently because the neutrino distribution functions are unknown. In the following paragraph, we will discuss the assumptions made for the evolution of the electron fraction in the neutrino driven wind which go back to Qian and Woosley (1996).

Applying the theory of weak interactions based on the reaction rates λ_{ij} for the reaction particles (i, j), i.e. electron and positron as well as electron neutrino and antineutrino captures, the evolution equation for the electron and positron fractions can be written as follows

$$\frac{dY_{e^-}}{dt} = -\lambda_{e^-p}Y_{e^-}Y_p + \lambda_{\nu_e n}Y_{\nu_e}Y_n, \quad (16)$$

$$\frac{dY_{e^+}}{dt} = -\lambda_{e^+n}Y_{e^+}Y_n + \lambda_{\bar{\nu}_e p}Y_{\bar{\nu}_e}Y_p. \quad (17)$$

These expressions can be combined to calculate the evolution of the total number of charges, using the relations $Y_p = Y_e$ and $Y_n = 1 - Y_e$,

$$\begin{aligned} \frac{dY_e}{dt} &= \frac{d}{dt}(Y_{e^-} - Y_{e^+}) \\ &= \lambda_{e^+n}Y_{e^+} + \lambda_{\nu_e n}Y_{\nu_e} \\ &\quad - (\lambda_{e^-p}Y_{e^-} + \lambda_{e^+n}Y_{e^+} + \lambda_{\nu_e n}Y_{\nu_e} + \lambda_{\bar{\nu}_e p}Y_{\bar{\nu}_e})Y_e, \end{aligned}$$

assuming fully dissociated nuclear matter. This expression is approximated in a crucial but powerful way (Qian and Woosley (1996) Eq.(73)), ignoring contributions from electron and positron captures as well as the decoupling of radiation and the angular dependency of the neutrino distribution function on the distance from the energy-dependent neutrinospheres, as follows


$$Y_e \simeq \frac{\lambda_{e^+n} Y_{e^+} + \lambda_{\nu_e n} Y_{\nu_e}}{\lambda_{e^-p} Y_{e^-} + \lambda_{e^+n} Y_{e^+} + \lambda_{\nu_e n} Y_{\nu_e} + \lambda_{\bar{\nu}_e p} Y_{\bar{\nu}_e}} \quad (18)$$

$$\simeq \frac{\lambda_{\nu_e n} Y_{\nu_e}}{\lambda_{\nu_e n} Y_{\nu_e} + \lambda_{\bar{\nu}_e p} Y_{\bar{\nu}_e}}. \quad (19)$$

This approximation of the electron fraction was further simplified and expressed in terms of the neutrino luminosities L_ν and mean neutrino energies $\langle E_\nu \rangle$ and the well known rest mass difference between neutron and proton $Q = m_n - m_p = 1.2935$ MeV, as follows

$$Y_e \simeq \left(1 + \frac{L_{\bar{\nu}_e} \langle E_{\bar{\nu}_e} \rangle - 2Q + \frac{1.2 Q^2}{\langle E_{\bar{\nu}_e} \rangle}}{L_{\nu_e} \langle E_{\nu_e} \rangle + 2Q + \frac{1.2 Q^2}{\langle E_{\nu_e} \rangle}} \right)^{-1}, \quad (20)$$

which is used in previous static steady-state and parametrized dynamic studies of the neutrino driven wind.



(a) Different electron fraction approximations (dash-dotted line: Eq. 19, dashed line: Eq. 20) in comparison with Boltzmann transport (solid line).

(b) Evolution of the electron fraction according to Eq. (20). The three lines correspond to different assumptions for the electron antineutrino mean energy.

Figure 13: The Electron fraction approximations at a distance of 10 km outside the electron-neutrinosphere for the $10.8 M_\odot$ progenitor model from Woosley et al. (2002).

Fig. 13(a) compares the electron fraction behavior at a distance of 10 km outside the electron-neutrinosphere, from Boltzmann neutrino transport (blue solid line) with the approximations Eq. (19) based on the neutrino capture rates (red dashed line) and Eq. (20) based on the luminosities and mean neutrino energies (red dash-dotted lines). The approximations are in qualitative agreement with Boltzmann transport. The differences on the longer timescale are most likely due to the presence of light and heavy nuclei which are not taken into account explicitly in the approximations. They change the number of free nucleons available for the reactions in Eq. (16) and (17). All descriptions agree qualitatively in the prediction of a generally proton-rich material in the wind, based on the neutrino spectra obtained via Boltzmann transport.

5.2 The neutrino observables in the wind

Comparing the neutrino spectra in Fig. 14 with the spectra assumed in previous static steady-state and dynamic wind studies (see for example Thompson et al. (2001) and Arcones et al. (2007)), we find two major differences: *One*, the neutrino luminosities and mean neutrino energies assumed are significantly larger than those we find and *two*, the assumed behaviour with respect to time is different.

The commonly used assumptions made in static steady-state and parametrized dynamic wind studies go back to the detailed investigation from Woosley et al. (1994), who performed core collapse simulations based on sophisticated input

physics. They investigated the neutrino driven explosion of a $20 M_{\odot}$ progenitor star and followed the evolution for 18 seconds post bounce into the neutrino driven wind phase. In their simulations the electron (anti)neutrino luminosities decreased from initially 4×10^{52} (3×10^{52}) erg/s at the onset of the explosion to 6×10^{51} (7.5×10^{50}) erg/s at 10 seconds after bounce, where strictly $L_{\bar{\nu}_e} > L_{\nu_e}$ after the onset of the explosion. The difference between the neutrino and antineutrino luminosities remained small and constant with respect to time up to 3 seconds post bounce and increased only significantly after 4 – 5 seconds post bounce, after which the difference reached its maximum of 1.5×10^{50} erg/s at the end of the simulation at about 18 seconds post bounce. The electron flavor neutrino luminosities in our models follow a different behaviour. They reach 1×10^{51} erg/s at about 5, 6 and 8 seconds post bounce for the 8.8, 10.8 and $18 M_{\odot}$ progenitor models respectively. The larger electron flavor neutrino luminosities for the more massive progenitors are in correlation with the more massive PNSs and the hence larger number of neutrinos emitted. However, the difference between electron-neutrino and electron-antineutrino luminosities found in the present investigation is significantly smaller than the difference in Woosley et al. (1994). During the initial explosion phase until about 300 ms after the onset of the explosion, the electron antineutrino luminosity is slightly larger than the electron neutrino luminosity by about 1×10^{50} erg/s which explains the electron fraction of $Y_e > 0.5$ of the early explosion ejecta. After about 900 ms post bounce, the luminosities can hardly be distinguished where during the initial neutrino driven wind phase after about 1 second after bounce the electron neutrino luminosity becomes larger than the electron antineutrino luminosity by about 1×10^{50} erg/s. This difference reduces again at later times at about 6 seconds post bounce and the electron flavor neutrino luminosities become more and more similar (see Fig. 14).

Even more different are the values and the behavior of the mean neutrino energies, see Fig. 14 and compare with Fig. 2 of Woosley et al. (1994). They found (μ/τ) -neutrino energies of about 35 MeV which remained constant with respect to time. Their electron-antineutrino energies increased slightly from about 20 MeV to 22 MeV where the electron-neutrino energies decrease from 14 MeV to 12 MeV. This increasing difference between the electron neutrino and antineutrino spectra favored neutron-rich material, which was consistent with their findings of $Y_e < 0.5$ for the material ejected in the neutrino driven wind in Woosley et al. (1994). We cannot confirm these results for the mean neutrino energies nor the evolution of the spectra. In contrast, all mean neutrino energies decrease with respect to time for all our models. This is a consequence of lepton number and energy loss of the central PNS where the neutrinos diffuse out. The electron (anti)neutrino energies decrease from about 10 (12) MeV at the onset of the explosion to about 8.5 (9) MeV and the (μ/τ) -neutrino energies decrease from 16 MeV to 10 MeV at the end of the simulations. Hence, not only the mean energies decrease also the difference between the electron flavor neutrino spectra decreases. The reason for the neutrino spectra to become more similar with respect to time is related to the evolution of the thermodynamic properties at the neutrinospheres, and will be discussed in the following subsection.

5.3 The PNS contraction

The behaviour of the neutrino spectra and hence the evolution and the properties of the neutrinospheres is related to the PNS contraction. The contraction is caused by a continuous deleptonization and translates to a continued steepening of the density gradient at the PNS surface. Hence, the neutrinosphere radii for the electron flavor neutrinos move closer together with time. The evolution of the neutrinosphere radii for both electron neutrino and antineutrino are illustrated in Fig. 15 (a) for the $10.8 M_{\odot}$ progenitor model. Their difference reduces from 740 m at about 1 second post bounce to 370 m at about 5 seconds post bounce and further to 260 m at about 10 seconds post bounce.

This contraction behaviour has consequences for the neutrino spectra, which are determined during the neutrino driven wind phase by diffusion rather than by mass accretion. Hence, the electron flavor neutrino luminosities can be determined as follows

$$L_{\nu} = \frac{1}{4} 4\pi r^2 u_{\nu} |_{R_{\nu}}, \quad (21)$$

where $u_{\nu} \propto T^4$ is the thermal black body spectrum for ultra-relativistic fermions with temperature T . The matter temperatures at the neutrinospheres decrease with respect to time as shown in Fig. 15 (b). This is due to the continued loss of lepton number and energy, carried away by the diffusing neutrino radiation field as illustrated in Fig. 16 for the $10.8 M_{\odot}$ from Woosley et al. (2002). The lepton number decreases from $Y_L \simeq 0.3$ at 2 seconds post bounce to $Y_L \simeq 0.18$ at 10 seconds post bounce, see Fig. 16 (c). The additionally reduced mean neutrino energies (on average), from $\langle E \rangle_{\text{rms}} \simeq 150$ MeV to $\langle E \rangle_{\text{rms}} \simeq 50$ MeV (see Fig. 16 (d)), and the consequent reduced temperature-peak inside the PNS (see Fig. 16 (b)) cause the contraction of the outer layers of the PNS. This can be identified via the density




Figure 14: Neutrino luminosities and mean energies with respect to time after bounce for the 8.8 $M_{\odot}$ O-Ne-Mg-core from Nomoto (1983,1984,1987) (left panel) and the 10.8 $M_{\odot}$ (middle panels) and 18 $M_{\odot}$ (right panel) Fe-core progenitor models from Woosley et al. (2002), measured in a co-moving reference frame at a distance of 500 km.

increase in Fig. 16 (a). Note in addition to the maximum temperature decrease at the outer layers of the PNS, the central temperature increases from 18 MeV to 23 MeV on the post bounce times between 2 and 10 seconds. This is caused by the contraction of the deleptonizing outer layers of the PNS which compresses the central part. The evolution of the reducing temperature at the neutrinospheres is shown in Fig. 15 (b). In combination with the loss of leptons number, it explains the decreasing electron flavor neutrino luminosities and mean neutrino energies with respect to time. Furthermore, the temperature difference decreases with respect to time from 0.467 MeV at about 1 second post bounce to 0.362 MeV at about 10 seconds post bounce. Consequently the neutrino spectra become more similar with respect to time. As illustrated in Fig. 14, the difference in the electron flavor neutrino luminosities and mean neutrino energies decreases for all models under investigation. In addition, Fig. 15 (c) and (d) illustrate the evolution of the baryon density and the electron fraction at the corresponding neutrinospheres. It becomes additionally clear from the electron fraction approximation Eq.(20), that it is not the absolute values for the mean neutrino and antineutrino energies that determine whether matter becomes neutron- or proton-rich but their ratio.




Figure 15: Evolution of the neutrinosphere radii in graph (a) and temperature and density at the corresponding neutrinospheres in graphs (b) and (c) respectively for the $10.8 M_{\odot}$ progenitor model from Woosley et al. (2002).




Figure 16: Radial PNS profiles at two different post bounce times (thick lines: 2 seconds, thin lines: 10 seconds), for the $10.8 M_{\odot}$ progenitor model from Woosley et al. (2002).

Since this difference is small in our simulations, with initially $\langle E_{\nu_e} \rangle_{\text{rms}} \simeq 10$ MeV and $\langle E_{\bar{\nu}_e} \rangle_{\text{rms}} \simeq 13$ MeV (at about 1 second post bounce) and only $\langle E_{\nu_e} \rangle_{\text{rms}} \simeq 9$ MeV and $\langle E_{\bar{\nu}_e} \rangle_{\text{rms}} \simeq 11$ MeV (at later times at 10 seconds post bounce), the values found for the electron fraction of $Y_e > 0.5$ (solid line in Fig. 13(b) for the Y_e -approximation based on the luminosity and mean neutrino energies) clearly illustrate that the accelerated matter in the neutrino driven wind stays proton-rich for more than 10 seconds. This is in qualitative agreement with Boltzmann transport as discussed above and shown in Fig. 13(a). Hence we find Eq.(20) to be a good approximation to model the electron fraction in the wind. On the other hand, most of the previous studies select the neutrino luminosities and mean energies to investigate a neutron-rich neutrino driven wind. In order to test the appearance of $Y_e < 0.5$ under such conditions, we increase the difference between the mean neutrino and antineutrino energies by hand. We evaluate expression (20) shown in

Fig. 13(b) at 10 km outside the electron-neutrinosphere for 1.2 (dashed line) and 1.5 (dash-dotted line) times larger electron-antineutrino mean energies. For the first value, Y_e decreases but matter remains slightly proton-rich, which for the latter value matter becomes neutron-rich. Indeed, the larger the difference between neutrino and antineutrino spectra are, the smaller becomes the electron fraction in the wind. Note that the luminosities and electron-neutrino energies remained unmodified for this experiment. Such an increase of the energy difference between neutrinos and antineutrinos could perhaps be related to the uncertainty of the EoS for nuclear matter, which will be discussed in the following paragraph.

The assumed PNS radii in previous wind studies reach about 10 km shortly (≤ 1 second) after the explosion has been launched. We define the radius of the PNS as the position of the electron-neutrinosphere at the steep density gradient at the PNS surface. The approximated inner boundary of the physical domain in most wind models is close to but still inside this radius. The position of the neutrinospheres and the contraction of the PNSs found in the present paper differ significantly from the assumptions made in most previous wind studies. We find PNS radii of about 40 km at the time of the explosion and 20 km at about 2 seconds after bounce. During the later evolution, the PNS contraction slows down. The PNS profile and hence the position of the neutrinospheres as well as the contraction behavior itself is given implicitly by the EoS for hot and dense nuclear matter as well as the PNS deleptonization. For the stiff EoS from Shen et al. (1998) and both the 10.8 and 18 $M_\odot$ progenitors, the PNSs reach radii of 14.5 – 15 km only at about 10 seconds after bounce (see Fig. 15). The larger radii of the neutrinospheres result in lower neutrino luminosities and mean energies and a smaller difference between neutrino and antineutrino spectra in comparison to the assumptions made in most previous wind models. This is in agreement with Arcones et al. (2007) who additionally assume PNS radii of 15 km and find conditions that differ more from previous wind studies. They obtained significantly larger values for the electron fraction. To summarize, this effects and the different behaviour of the neutrino spectra assumed in the previous wind studies leads to different matter properties of the neutrino driven wind. A detailed comparison study of fast and slowly contracting PNSs with respect to the neutrino driven wind, e.g. applying EoSs with different compressibilities and asymmetry energies, would be necessary in the context of radiation hydrodynamics simulations using spectral Boltzmann neutrino transport.

6 Long term post bounce evolution

During the neutrino driven wind phase, the neutrino luminosities and mean neutrino energies decrease continuously, which leads to a constant decrease in the net-heating rates. At luminosities below 10^{51} erg/s (see Fig. 14), the supersonic matter outflow for the 10.8 $M_\odot$ progenitor model descends into a subsonic expansion. The wind termination shock turns again into a subsonic neutrino driven wind. At later times, the neutrino driven wind settles down to a quasi-stationary state with no significant matter outflow, illustrated at the example of the 18 $M_\odot$ progenitor model in Fig.17 (a). The explosion shock continues to expand and the material enclosed inside the mass cut accretes onto the PNS at the center. In combination with the deleptonization, this leads to the continuous PNS contraction. However, the contraction proceeds on a timescale of seconds and hence the PNS can be considered in a quasi-stationary state. The dense and still hot and lepton-rich PNS at the center is surrounded by a low density and high entropy atmosphere, composed of light and heavy nuclei. See for example the abundances of the 18 $M_\odot$ progenitor for the post bounce time of 22 seconds in Fig. 17 (e).

The internal temperature profile of the PNS is not constant. The central region of the PNS did not experience shock heating immediately after the Fe-core bounce, since the initial shock forms at the edge of the bouncing core. Its mass scales roughly with Y_e^2 and is typically around values of 0.5 – 0.6 $M_\odot$ for low and intermediate mass Fe-core progenitors. Hence, the central temperature after bounce is given by the thermodynamic conditions at bounce. The temperature changes only during the post bounce evolution due to compressional heating and the diffusion of neutrinos. The shock heated material inside the PNS shows significantly higher temperatures than at the center. The temperature decreases again towards the PNS surface where the matter is less dense (for the illustration of the radial temperature profile inside the PNS as well as the dynamical evolution of temperature and density, see Fig.17 (f) and (b) at selected post bounce times between 5 – 22 seconds). The neutrinos diffuse continuously out of the PNS and carry away energy. The central electron fraction reduces from $Y_e \approx 0.25$ at the onset of the explosion to $Y_e \approx 0.15$ at 22 seconds after bounce (see Fig.17 (d)). It relates to a temperature decrease from about 35 MeV initially (at 3 seconds post bounce) to 23 MeV at about 22 seconds post bounce. This corresponds to the initial and neutrino dominated cooling phase.




Figure 17: Radial profiles of selected hydrodynamic variables for the 18 $M_{\odot}$ progenitor model at three different post bounce times, illustrating the disappearance of the neutrino driven wind and the PNS cooling and contraction. Graph (e) illustrates the composition at 22 seconds post bounce.

Unfortunately the achieved temperatures are not representative since important neutrino reactions, such as the direct and modified Urca processes, are not yet taken into account.




Figure 18: Comparing selected hydrodynamic variables using the standard reactions rates based on Bruenn (1985) (solid lines) and the artificially enhanced rates (dashed lines) for the 10.8 M_⊙ (top) and the 18 M_⊙ progenitor model (bottom).

7 Discussion

The neutrino driven wind was found to occur in all three progenitor models under investigation, the 8.8 M_⊙ O-Ne-Mg-core and the 10.8 and 18 M_⊙ Fe-core progenitor models. Because the neutrino driven explosions for the Fe-core progenitors are launched using artificially enhanced neutrino reaction rates, one may ask about the impact of these modified rates on the neutrino driven wind. Therefore we performed additional runs for which we switch back to the standard opacities given in Bruenn (1985) after the explosion has been launched. The times when we switch back is about 500 ms after bounce, chosen such that the dynamics of the explosion ejecta does not change anymore significantly due to neutrino heating. However, the lower opacities translate to a significantly smaller net-heating by a factor of 5 – 6 in the region on top of the PNS where the neutrino driven wind develops. The energy deposition is still sufficient to drive the neutrino driven wind but the matter velocities are smaller by a factor of 2 – 5 in comparison to the wind velocities using the enhanced reaction rates (see Fig. 18 (a)). The main effect of the artificially enhanced reaction rates and the hence increased neutrino heating on the dynamics is clearly the stronger neutrino driven wind. For the 10.8 M_⊙ progenitor model and with the enhanced heating, the wind even develops supersonic velocities (as discussed above in § 4) in Fig. 18 (a) (top panel). The supersonic wind collides with the explosion ejecta where matter decelerates and hence the reverse shock forms, which additionally increases the entropy in the wind (see Fig. 18(c) (top panel)). This additional entropy increase is absent in the simulations using the standard opacities, where the wind stays subsonic. This is also the case for the 18 M_⊙ progenitor model (Fig. 18, bottom panel), with and without the enhanced opacities. The neutrino driven wind of the O-Ne-Mg-core is illustrated in Fig. 11 using the standard rates based on Bruenn (1985). The formation of a supersonic neutrino driven wind could be confirmed including the formation of the wind termination shock. Hence, one may speculate whether only low mass progenitors develop a strong neutrino driven outflow where the influence of the wind to the matter properties of the ejecta becomes small

for more massive progenitors. The progenitor dependency of the neutrino driven wind is related to the density of the envelope surrounding the PNS after the explosion has been launched, which is significantly higher for more massive progenitors and hence the neutrino driven wind is weaker.

The agreement of the time evolution of the mean neutrino energies between all three progenitor models under investigation (using the enhanced and standard opacities) in Fig. 14 is striking. The impact of the artificial heating to the neutrino observables and hence to the electron fraction in the wind is less pronounced. The influence on the composition of the wind is illustrated via the electron fraction in Fig. 18 (d). Using the standard rates, the wind stays slightly more proton-rich. Increasing the charged current reaction rates allows β -equilibrium to be established on a shorter timescale. In addition, matter stays slightly more proton-rich for the the less intense neutrino driven wind, which develops for the Fe-core progenitors using the standard neutrino opacities. The additional electron fraction reduction in the neutrino driven wind for the models using the enhanced neutrino reactions is found due to the larger degeneracy obtained in the stronger deceleration behind the explosion ejecta. However, the findings of generally proton-rich ejecta as well as the generally proton-rich neutrino driven wind does not change. The corresponding densities and entropies per baryon in the wind are shown in Fig. 18 (b) and (c). The effects of the artificial heating are slightly smaller entropies. The higher matter outflow velocities in the wind region using the artificial heating results additionally in lower densities, shown in Fig. 18 (b).

The artificially increased charged current reaction rates cannot be justified by physical uncertainties of the rates themselves. Similar to the large luminosities assumed in Arcones et al. (2007), they could rather be seen as a lowest order attempt to take the effects of multi-dimensional phenomena into account. For example, known fluid instabilities increase the neutrino energy deposition efficiency. Convection allows matter to stay for a longer time in the neutrino heating region (see Herant et al. (1994), Janka and Müller (1996)). Present axially symmetric core collapse models of massive Fe-core progenitor stars (even non-rotating) predict bipolar explosions (see Janka et al. (2008)). The deviation from a spherical description and hence the deformation of the SAS due to fluid instabilities takes place during the neutrino heating phase on a timescale of several 100 ms after bounce. In multi-dimensional models, the luminosities are powered by a significantly larger mass accretion because the up-streaming neutrino heated matter is accompanied by a large down-stream of cold material. These larger luminosities may power a supersonic neutrino driven wind behind the explosion ejecta while the neutrino driven wind may remain absent in the angular directions of the accreting material which will not be ejected.

Previous wind models have long been investigated as a possible site for the production of heavy elements via the r -process, motivated by the expectation of the ejection of neutron rich material, the large entropies in the neutrino driven wind and the short timescale of the neutrino driven wind expansion (see Hoffman et al. (2007) and Panov and Janka (2009) and references therein). The relevant quantities are shown in Figs. 10 and 11 for the $10.8 M_{\odot}$ and $8.8 M_{\odot}$ progenitor models respectively. Illustrated are several selected mass shells that are part of the region where the neutrino driven wind develops in our radiation hydrodynamics model based on spectral three flavor Boltzmann neutrino transport. The inclusion of neutrino transport in a dynamical model is essential in order to obtain consistent neutrino spectra which determine the evolution of the electron fraction and the PNS contraction due to deleptonization and mass accretion. In comparison to previous static steady-state and dynamic wind models - where these ingredients were assumed - we confirm several properties of the accelerated material in the neutrino driven wind, such as the fast expansion and the large matter outflow rate shown in Fig. 19(a) and (b), the high velocities in the Figs. 10(f) and 11(f) and the rapidly decreasing density and temperature of the accelerated material in Fig. 10 (b) and (c) respectively. The expansion timescale in Fig. 19(b) is given by the following expression

$$\tau_{\text{dyn}} = \left. \frac{r}{v} \right|_{T=0.5 \text{ MeV}},$$

evaluated at the surface of constant temperature of $T = 0.5 \text{ MeV}$, compared with an alternative definition of the dynamic timescale which has been introduced in Thompson et al. (2001)

$$\tau_{\rho} = \left| \frac{1}{v} \frac{1}{\rho} \frac{\partial \rho}{\partial r} \right|^{-1}_{T=0.5 \text{ MeV}},$$

as well as a timescale approximation which has been derived in Qian and Woosley (1996) Eq.(61)

$$\tau_{\text{QW}} \propto \frac{1}{L_{\bar{\nu}_e}} \frac{1}{\epsilon_{\bar{\nu}_e}} R_{\text{PNS}} M_{\text{PNS}},$$

where additionally an approximation for the mass outflow rate is derived as follows

$$\left. \frac{dM}{dt} \right|_{\text{QW}} \propto L_{\bar{\nu}_e}^{5/3} \epsilon_{\bar{\nu}_e}^{10/3} R_{\text{PNS}}^{5/3} M_{\text{PNS}}^{-2},$$

where R_{PNS} and M_{PNS} are the PNS radius and mass respectively, which we take to be given by the electron-antineutrinosphere, and $\epsilon_{\bar{\nu}_e} = \langle E_{\bar{\nu}_e}^2 \rangle / \langle E_{\bar{\nu}_e} \rangle$ where $\langle E_{\bar{\nu}_e} \rangle$ is the mean electron-antineutrino energy and $L_{\bar{\nu}_e}$ is the electron-antineutrino luminosity, both taken at the surface of constant temperature of $T = 0.5$ MeV. In comparison with previous wind studies (see for example Fig. 4 of Arcones et al. (2007)), we find generally a longer timescale of $\tau_{\text{dyn}} = 30 - 100$ ms for the neutrino driven wind which corresponds to a mass outflow rate of $10^{-3} - 10^{-5} M_{\odot} \text{ s}^{-1}$ shown in Fig. 19 (a) and (b) respectively. However, we find generally a much longer timescale than the approximation τ_{QW} which is of the order $\sim 1 - 5$ ms. The same holds for the mass outflow rate $dM/dt|_{\text{QW}}$ which is smaller by one order of magnitude. This may be related to the crucial assumptions made during the derivation of the above expressions, e.g. hydrostatic equilibrium, $R_{\text{PNS}} = 10$ km, $L_{\bar{\nu}_e} \simeq L_{\nu_e}$, $\epsilon_{\bar{\nu}_e} = 20$ MeV, which differ significantly from our findings. However, the wind entropies of $40 - 100 k_B$ found (initially driven due to neutrino heating and additionally due to the deceleration in the reverse shock) are significantly smaller than often assumed in the literature and the previously accelerated matter does not become neutron rich as the neutrino driven wind decelerates behind the explosion ejecta but stays slightly proton-rich where $Y_e \simeq 0.54$ for more than 10 seconds. This, in combination with the much slower PNS contraction illustrated via the neutrinospheres in Fig. 15 in comparison to static steady-state and dynamic wind models suggest that the assumptions made in previous wind studies should be carefully reconsidered. With respect to Woosley et al. (1994) (e.g. Fig. 3), we find generally smaller mean neutrino energies which decrease with respect to time after bounce. This results in a decreasing difference between the electron flavor neutrino mean energies, while in Woosley et al. (1994) this difference increases. This fact in combination with the different PNS properties found in Woosley et al. (1994), enabled a strong neutrino driven wind where high entropies up to $400 k_B/\text{baryon}$ and a low electron fraction of $Y_e \simeq 0.35 - 0.45$ was obtained. These properties of the neutrino driven wind differ quantitatively from our results, where only entropies of $60 - 100 k_B/\text{baryon}$ are obtained and matter stays proton-rich ($Y_e \simeq 0.52 - 0.54$) for more than 10 seconds.




Figure 19: Comparing the mass accretion rate from our simulations and the approximation from Qian and Woosley (1996) in the graphs (a) and different dynamic timescale in the graphs (b) with respect to time after bounce for the 10.8 (left panel) and $18 M_{\odot}$ (right panel) progenitor models.

8 Summary and Outlook

For the first time, spherically symmetric core collapse supernova simulations based on general relativistic radiation hydrodynamics and three flavor Boltzmann neutrino transport are performed consistently for more than 20 seconds. We follow the dynamical evolution of low and intermediate mass progenitors through the collapse, bounce, post bounce, explosion and neutrino driven wind phases. The explosions of Fe-core progenitors of 10.8 and $18 M_{\odot}$ are modeled using artificially enhanced opacities, while the explosion of the $8.8 M_{\odot}$ O-Ne-Mg-core was obtained using the standard opacities. For all models under investigation, we confirm the formation and illustrate the conditions for the appearance of the neutrino driven wind during the dynamical evolution after the explosions have been launched. For the O-Ne-Mg-core and the $10.8 M_{\odot}$ Fe-core progenitor models, the supersonic neutrino driven wind collides with the slower

expanding explosion ejecta where due to the deceleration a reverse shock forms, known as the neutrino driven wind termination shock. We discuss the impact of the reverse shock on several properties of the ejecta and find general agreement with Arcones et al. (2007).

The comparison with approximate and static steady-state as well as parametrized dynamic wind models leads to a discrepancy in the obtained physical properties of the neutrino driven wind. Although the evolution of the hydrodynamic variables are in general agreement, we find smaller neutrino luminosities and a different behavior of the mean neutrino energies. In particular, the differences between the neutrino and antineutrino luminosities and mean neutrino energies are smaller. These differences reduce with time as the PNS contracts and result in a generally proton-rich neutrino driven wind over more than 10 seconds. Hence, the suggestion that the physical conditions in the neutrino driven wind could be favorable for the nucleosynthesis of heavy elements via the r -process could not be confirmed. For a more accurate determination of the yields of the neutrino driven wind, a detailed nucleosynthesis analysis taking the r -, p - and νp -processes into account is required. In order to further support the robustness, improvements of the input physics such as weak magnetism and nucleon-nucleon recoil (following e.g. Horowitz (2002)), taking the presence of light and heavy clusters of nuclei into account as well as different EoSs with respect to the different PNS contraction behavior, should be considered. These may have a strong influence on the properties of the neutrino spectra at the neutrinospheres and may therefore modify some of the results found in the present study of the neutrino driven wind.

Our simulations are carried out until the neutrino driven wind settles down to a quasi-stationary state leading to the initial and neutrino dominated PNS cooling phase. There, the simulations have to be stopped because important neutrino cooling processes like the direct and modified URCA processes are not taken into account yet. However, a smooth connection to isolated neutron or protoneutron star cooling studies comes into reach for future work (Henderson and Page (2007)).

Acknowledgment

The authors would like to thank A. Arcones and G. Martínez-Pinedo for discussions and helpful comments. The project was funded by the Swiss National Science Foundation grant numbers PP002-106627/1, PP00P2-124879 and 200020-122287. The authors are additionally supported by CompStar, a research networking program of the European Science Foundation, and the Scopes project funded by the Swiss National Science Foundation grant. no. IB7320-110996/1. A. Mezzacappa is supported at the Oak Ridge National Laboratory, which is managed by UT-Battelle, LLC for the U.S. Department of Energy under contract DE-AC05-00OR22725.

References

- Arcones, A.; Janka, H.-Th.; and Scheck L. 2007, *A&A*, 467, 1227
- Arcones, A.; Martínez-Pinedo, G.; O’Conner, E.; Schwenk, A.; Janka, H.-Th.; Horowitz, C. J.; and Langanke, K. 2008, *PRC*, 78, 015806-+
- Bethe, H. A. and Wilson, J. R. 1985, *ApJ*, 295, 14
- Bruenn, S. W. 1985, *ApJS*, 58,771
- Bruenn, S. W.; Dirk, C. J.; Mezzacappa, A.; Hayes, J. C.; Blondin, J. M.; Hix, W. R. and Messer, O. E. B. 2006, *Journal of Physics Conference Series*, 46, 393
- Burrows, A.; Hayes, J. and Fryxell, B. A. 1995, *ApJ*, 450, 830
- Duncan, R. C.; Shapiro, S. L. and Wasserman, I. 1986, *ApJ*, 309, 141
- Fischer, T.; Whitehouse, S. C.; Mezzacappa, A.; Thielemann, F.-K. and Liebendörfer, M. 2009, *A&A*, 499, 1
- J. L. Fisker. 2004, Ph.D. thesis, private communications

- Fröhlich, C.; Hauser, P.; Liebendörfer, M.; Martínez-Pinedo, G.; Thielemann, F.-K.; Bravo, E.; Zinner, N. T.; Hix, W. R.; Langanke, K.; Mezzacappa, A. and Nomoto, K. 2006a *ApJ*, 637, 415
- Fröhlich, C.; Hix, W. R.; Martínez-Pinedo, G.; Liebendörfer, M.; Thielemann, F.-K.; Bravo, E.; Langanke, K. and Zinner, N. T. 2006b, *New Astronomy Review*, 50, 496
- Fröhlich, C.; Martínez-Pinedo, G.; Liebendörfer, M.; Thielemann, F.-K.; Bravo, E.; Hix, W.-R.; Langanke, K. and Zinner, N. T. 2006c *Physical Review Letters*, 96 (14), 142502
- Gögelein, P. 2007, private communications
- Henderson, J. A. and Page, D. 2007, *Ap&SS*, 308, 513
- Herant, M.; Benz, W.; Hix, W. R.; Fryer, C. L. and Colgate, S. A. 1994, *ApJ*, 435, 339
- Hix, W. R.; Messer, O. B. E.; Mezzacappa, A.; Liebendörfer, M.; Sampaio, J.; Langanke, K.; Dean, D. J. and Martínez-Pinedo, G. 2003, *Physical Review Letters*, 91, 201102
- Hoffman, R. D.; Pruet, J.; Fisker, J. L.; Janka, H.-Th.; Buras, R. and Woosley, S. E. 2007, *ArXiv e-prints* 0712.2847
- Hoffman, R. D.; Woosley, S. E. and Qian, Y. Z. 1997, *ApJ*, 482, 951
- Horowitz, C. J. 2002, *PRD*, 65, 043001
- Janka, H.-Th. and Müller, E. 1995, *ApJL*, 448, L109
- Janka, H.-Th. and Müller, E. 1995, *A&A*, 306, 167
- Janka, H.-Th. 2001, *A&A*, 368, 527
- Janka, H.-Th.; Buras, R.; Kitaura Joyanes, F. S.; Marek, A.; Rampp, M. and Scheck, L. 2005, *Nuclear Physics A*, 758, 19
- Janka, H.-Th.; Marek, A.; Müller, B. and Scheck, L. 2008, In C. Bassa, Z. Wang, A. Cumming, and V. M. Kaspi, editors, *40 Years of Pulsars: Millisecond Pulsars, Magnetars and More*, volume 983 of *American Institute of Physics Conference Series*, pages 369–378
- Kitaura, F. S.; Janka, H.-Th. and Hillebrandt, W. 2006, *A&A*, 450, 345
- Kotake, K.; Sato, K. and Takahashi, K. 2006, *Reports on Progress in Physics*, 69, 971
- Langanke, K.; Martínez-Pinedo, G.; Sampaio, J. M.; Dean, D. J.; Hix, W. R.; Messer, O. B. E.; Mezzacappa, A.; Liebendörfer, M.; Janka, H.-Th. and Rampp, M. 2003, *Physical Review Letters*, 90, 241102
- Liebendörfer, M.; Mezzacappa, A.; Thielemann, F.-K.; Messer, O. E. B.; Hix, W. R. and Bruenn, S. 2001a, *PRD*, 63, 103004
- Liebendörfer, M.; Mezzacappa, A. and Thielemann, F.-K. 2001b, *PRD*, 63, 104003
- Liebendörfer, M.; Rosswog, S. and Thielemann, F.-K. 2002, *ApJS*, 141, 229
- Liebendörfer, M. 2004, *Proceedings of the 12th Workshop on Nuclear Astrophysics*, Rep. No. MPA/P14, 143, *eprint*, astro-ph/0405029
- Liebendörfer, M.; Messer, O. E. B.; Mezzacappa, A.; Bruenn, S. W.; Cardall, C. Y. and Thielemann, F.-K. 2004, *ApJS*, 150, 263
- Liebendörfer, M.; Rampp, M.; Janka, H.-Th. and Mezzacappa, A. 2005, *ApJ*, 620, 840
- Marek, A. and Janka, H.-Th. 2009, *ApJ*, 694, 664

- Mayle, R. and Wilson, J. R. 1988, *ApJ*, 334, 909
- Messer, O. E. B. and Bruenn, S. W. 2003, private communications
- Mezzacappa, A.; Blondin, J. M.; Messer, O. E. B. and Bruenn, S. W. 2006, In *Origin of Matter and Evolution of Galaxies*, volume 847 of *American Institute of Physics Conference Series*, pages 179–189
- Mezzacappa, A. and Bruenn, S. W. 1993a, *ApJ*, 405, 637
- Mezzacappa, A. and Bruenn, S. W. 1993b, *ApJ*, 405, 669
- Mezzacappa, A. and Bruenn, S. W. 1993c, *ApJ*, 410, 740
- Mezzacappa, A. and Messer, O. E. B. 1999, *Journal of Computational and Applied Mathematics*, 109, 281
- Miller, D. S.; Wilson, J. R. and Mayle, R. W. 1993, *ApJ*, 415, 278
- Misner, C.W. and Sharp, D. H. 1964, *Physical Review*, 136, 571
- Nomoto, K. 1983, *IAU Symposium*, 101, 139
- Nomoto, K. 1984, *ApJ*, 277, 791
- Nomoto, K. 1987, *ApJ*, 322, 206
- Otsuki, K.; Tagoshi, H.; Kajino, T. and Wanajo, S. Y. 2000, *ApJ*, 533, 424
- Panov, I. V. and Janka, H.-Th. 2009, *A&A*, 494, 829
- Qian, Y. Z. and Woosley, S. E. 1996, *ApJ*, 471, 331
- Sagert, I.; Fischer, T.; Hempel, M.; Pagliara, G.; Schaffner-Bielich, J.; Mezzacappa, A.; Thielemann, F.-K. and Liebendörfer, M. 2009, *Physical Review Letters*, 102 (8) 081101
- Scheck, L.; Kifondis, K.; Janka, H.-T. and Müller, E. 2006, *A&A*, 457, 963
- Schinder, P. J. and Shapiro, S. L. 1982, *ApJS*, 50, 23
- Shen, H.; Toki, H.; Oyamatsu, K. and Sumiyoshi, K. 1989, *Progress of Theoretical Physics*, 100, 1013
- Takahashi, K.; Witt, J. and Janka, H.-Th. 1994, *A&A*, 286, 857
- Thielemann, F.-K.; Brachwitz, F.; Höflich, P.; Martinez-Pinedo, G. and Nomoto, K. 2004, *New Astronomy Review*, 48, 605
- Thompson, T. A. and Burrows, A. 2001, *Nuclear Physics A*, 688, 377
- Thompson, T. A.; Burrows, A. and Meyer, B. S. 2001, *ApJ*, 562, 887
- Timmes, F. X. and Arnett, D. 1999, *ApJS*, 125, 277
- Wanajo, S. 2006a, *ApJ*, 647, 1323
- Wanajo, S. 2006b, *ApJL*, 650, L79
- Wilson, J. R. and Mayle, W. R. 1993, *Phys. Rep.*, 227, 97
- Witt, J.; Janka, H.-Th. and Takahashi, K. 1994, *A&A*, 286, 841
- Woosley, S. E. and Baron, E. 1992, *ApJ*, 391, 228
- Woosley, S. E.; Wilson, J. R.; Mathews, G. J.; Hoffman, R. D. and Meyer, B. S. 1994, *ApJ*, 433, 229
- Woosley, S. E.; Heger, A. and Weaver, T. A. 2002, *Reviews of Modern Physics*, 74, 1015
- Yueh, W. R. and Buchler, J. R. 1976, *Ap&SS*, 41, 221